



Corrigendum to: 'Investigation of Cu heteroatoms and Cu clusters in Fe-Cu alloy and their special effect mechanisms on the Fenton-like catalytic activity and reusability' [Appl. Catal. B Environ. 299 (2021) 120662]

Qixing Xia ^{a,b,c,d}, Dongjie Zhang ^b, Zhongping Yao ^{b,*}, Zhaohua Jiang ^b

^a Institute of Culture and Heritage, Northwestern Polytechnical University, Xi'an 710000, China

^b School of Chemistry and Chemical Engineering, State Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology, Harbin 150001, China

^c School of Materials Science and Engineering, Northwestern Polytechnical University, Xi'an 710000, China

^d The Key Scientific Research Base of the State Administration of Cultural Relics for the Protection and Restoration of the Collection Murals and Materials Science Research, Xi'an 710000, China



The authors regret to mishandle the data in Fig. 8a when processing the original data. In the ESR tests, the curves of Fe-25Cu and ZVI at 1 min are very similar, and the authors unintentionally mixed the data, leading to the curve of Fe-25Cu replaced by that of ZVI. The corrected Fig. 8 is as

follow. The error does not change the overall results and discussion, as well as the final conclusion.

The authors would like to apologize for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.apcatb.2021.120662>.

* Corresponding author.

E-mail address: yaozhongping@hit.edu.cn (Z. Yao).




Fig. 8. The EPR spectra (a) of Fenton-like systems via ZVI and Fe-25Cu at 1 min and 10 min, and the optimized configurations (b) for H_2O_2 adsorption on the Fe atoms at the normal Fe lattice (down) and the Cu doped Fe lattice (up): Violet – Fe atom, reddish-Cu atom, white-H atom, red-O atom. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)